

Charged Heterotic Black-Holes in Four and Two Dimensions

Gabriel Lopes Cardoso ^a

Institute for Theoretical Physics, Utrecht University, 3508 TA Utrecht, Netherlands

ABSTRACT

We consider four-dimensional charged black-holes occuring in toroidally compactified heterotic string theory, whose ten-dimensional interpretation involves a Kaluza-Klein monopole and a five-brane. We show that these four-dimensional black-holes can be connected to two-dimensional charged heterotic black-holes upon removal of the constants appearing in the harmonic functions associated with the Kaluza-Klein monopole and the five-brane.

April 1998

^aEmail: cardoso@fys.ruu.nl

It was pointed out by Hyun in [1] that five-dimensional black-holes occuring in toroidally compactified type II string theory are related to two-dimensional black-holes by U-duality. In [2] it was shown that the associated entropies agree. Here, we shall consider a particular class of extremal four-dimensional black-holes occuring in toroidally compactified heterotic string theory and we will connect it to a class of extremal two-dimensional heterotic black-holes. The two-dimensional black-holes in question [3, 4] are derived from the heterotic string target-space action

$$\frac{1}{16\pi G_2} \int d^2x \sqrt{-g} e^{-2\phi} \left\{ R + 4(\nabla\phi)^2 - \frac{1}{4}F^2 + c \right\} . \quad (1)$$

We will show that the associated entropies match. It should be straightforward to extend this connection to the case of non-extremal black-holes.

The extremal four-dimensional heterotic black-holes we will consider are the ones obtained by compactifying the heterotic ten-dimensional solution describing the intersection of a string, a five-brane, a Kaluza-Klein monopole and a wave [5]. The ten-dimensional line element is given by

$$\begin{aligned} ds_{10}^2 = & \frac{1}{H_1} \left(-dt^2 + dy^2 + (H_0 - 1)(dt + dy)^2 \right) + H_5 H_{KK} dx_i^2 + \frac{H_5}{H_{KK}} (dz + A_i dx^i)^2 \\ & + dx_6^2 + dx_7^2 + dx_8^2 + dx_9^2 \quad , \quad r^2 = x_i^2 \quad , \quad i = 1, 2, 3 \end{aligned} \quad (2)$$

where the $H_i = 1 + \frac{Q_i}{r}$ ($i = 0, 1, 5, KK$) denote the harmonic functions associated with the wave, the string, the five-brane and the Kaluza-Klein monopole, respectively. Also $F_{ij} = \partial_i A_j - \partial_j A_i = \varepsilon_{ijk} \partial_k H_{KK}$. The ten-dimensional dilaton and the antisymmetric tensor are given by

$$e^{-2\phi} = \frac{H_1}{H_5} \quad , \quad B_{ty} = \frac{1}{H_1} \quad , \quad H_{zij} = \frac{1}{2} \varepsilon_{ijk} \partial_k H_5 \quad . \quad (3)$$

Dimensionally reducing in the y, z, x_6, x_7, x_8, x_9 directions and going to the Einstein frame yields the four-dimensional metric

$$ds_4^2 = -\frac{1}{\sqrt{H_0 H_1 H_5 H_{KK}}} dt^2 + \sqrt{H_0 H_1 H_5 H_{KK}} dx_i^2 \quad . \quad (4)$$

This describes an extremal four-dimensional black-hole, and its entropy is given by [6]

$$S = \frac{A}{4G_4} = \frac{\pi}{G_4} \sqrt{Q_0 Q_1 Q_5 Q_{KK}} \quad , \quad (5)$$

where A denotes the area of the horizon and G_4 denotes Newton's constant in four dimensions.

Before relating this four-dimensional black-hole to an extremal two-dimensional black-hole, let us first link the near-horizon geometry of the former to the geometry of a three-dimensional BTZ black-hole, as in [1, 7, 8]. This is achieved by dimensionally reducing (2)

in the z, x_6, x_7, x_8, x_9 directions and by going to the Einstein frame. In the near-horizon region ($Q_i/r \gg 1$) the resulting five-dimensional metric is given by

$$ds_5^2 = \frac{r}{(Q_1 Q_5 Q_{KK})^{\frac{1}{3}}} \left(-dt^2 + dy^2 + \frac{Q_0}{r} (dt + dy)^2 \right) + (Q_1 Q_5 Q_{KK})^{\frac{2}{3}} \left(\frac{dr^2}{r^2} + d\Omega_2^2 \right) . \quad (6)$$

The five-dimensional dilaton and the zz component of the metric are approximately constant and given by $e^{-2\phi} = Q_1/\sqrt{Q_5 Q_{KK}}$ and $g_{zz} = Q_5/Q_{KK}$, respectively. The five-dimensional metric can be rewritten as [8]

$$\begin{aligned} ds_5^2 &= ds_3^2 + \frac{l^2}{4} d\Omega_2^2 , \quad l = 2(Q_1 Q_5 Q_{KK})^{\frac{1}{3}} , \\ ds_3^2 &= -N^2 d\tau^2 + \frac{d\rho^2}{N^2} + \rho^2 (d\phi + N_\phi d\tau)^2 , \end{aligned} \quad (7)$$

where

$$N^2 = \frac{\rho^2}{l^2} - M + \left(\frac{8G_3 J}{2\rho} \right)^2 , \quad N_\phi = \frac{8G_3 J}{2\rho^2} , \quad M = \frac{8G_3 J}{l} = \frac{4Q_0}{l^3} , \quad (8)$$

and where $\tau = lt$, $\rho^2 = 2(r + Q_0)/l$, $\phi = y$, $0 \leq \phi \leq 2\pi$. This geometry describes a space which is the product of a two-sphere of radius $l/2$ and an extremal BTZ black-hole with mass M and angular momentum J . Note that the BTZ part and the sphere are completely decoupled. Inspection of (2) and (3) shows that the two-sphere supports a $U(1)$ gauge field. The extremal BTZ black-hole carries entropy [9]

$$S = \frac{\pi}{4G_3} \sqrt{2l^2 M} , \quad (9)$$

where G_3 denotes the three-dimensional Newton's constant. Using that the three and four-dimensional Newton's constants are related by [8] $\frac{1}{G_3} = \frac{1}{G_4} \frac{l^2}{2}$ it follows that the BTZ entropy (9) agrees with the macroscopic entropy (5) of the four-dimensional black-hole.

Next consider the five-dimensional metric in the string frame, obtained by reducing (2) over z, x_6, x_7, x_8, x_9 :

$$ds_5^2 = \frac{1}{H_1} \left(-dt^2 + dy^2 + (H_0 - 1)(dt + dy)^2 \right) + H_5 H_{KK} dx_i^2 , \quad g_{zz} = \frac{H_5}{H_{KK}} . \quad (10)$$

The five-dimensional dilaton reads $e^{-2\phi} = H_1/\sqrt{H_5 H_{KK}}$. Let us now drop the constant part in the harmonic functions H_5 and H_{KK} associated with the five-brane and the Kaluza-Klein monopole, so that the resulting space-time is no longer asymptotically flat. Below we will argue that the constant part in these harmonic functions should be removable by a sequence of S and T - S - T duality transformations in four dimensions. Let us furthermore identify $H_0 = H_1$. Then, the line element (10) turns into

$$ds_5^2 = \frac{1}{H_1} \left(-dt^2 + dy^2 + (H_1 - 1)(dt + dy)^2 \right) + Q_5 Q_{KK} \left(\frac{dr^2}{r^2} + d\Omega_2^2 \right) . \quad (11)$$

Note that $g_{zz} = Q_5/Q_{KK}$ is now constant. Dimensionally reducing (11) in the y direction yields

$$ds_4^2 = -\frac{1}{H_1^2} dt^2 + Q_5 Q_{KK} \frac{dr^2}{r^2} + Q_5 Q_{KK} d\Omega_2^2 \quad , \quad e^{-2\phi} = \frac{r + Q_1}{\sqrt{Q_5 Q_{KK}}} \quad . \quad g_{yy} = 1 \quad . \quad (12)$$

This metric describes the product of two two-dimensional spaces, one of which we will relate to a two-dimensional black-hole and one which is a two-sphere with radius $\sqrt{Q_5 Q_{KK}}$. Inspection of (2) and (3) shows that each of the two-dimensional geometries supports a $U(1)$ gauge field. Both two-dimensional parts are completely decoupled. Note that since the internal components of the metric g_{yy} and g_{zz} are constant, the only non-trivial scalar field is the dilaton.

The solution (12) can now be related to the two-dimensional heterotic black-hole solution of McGuigan, Nappi and Yost [3] as follows. The latter is described by

$$\begin{aligned} ds_2^2 &= -(1 - 2me^{-Qx} + q^2 e^{-2Qx}) dt^2 + (1 - 2me^{-Qx} + q^2 e^{-2Qx})^{-1} dx^2 \quad , \\ e^{-2(\phi-\phi_0)} &= e^{Qx} \quad , \quad F_{tx} = \sqrt{2} Q q e^{-Qx} \quad . \end{aligned} \quad (13)$$

Here m and q are constants related to the mass and to the electric charge of the solution, with $m > 0$ and $m^2 \geq q^2$. Q is a positive constant which determines the central charge deficit $c = Q^2$ appearing in the action (1). The asymptotic flat region corresponds to $x = \infty$, whereas the curvature singularity is at $x = -\infty$. By changing the spatial variable to $y = e^{-Qx}$, the solution (13) becomes

$$\begin{aligned} ds_2^2 &= -q^2 (y - y_1)(y - y_2) dt^2 + \frac{1}{q^2 Q^2} \frac{dy^2}{y^2 (y - y_1)(y - y_2)} \quad , \\ e^{-2(\phi-\phi_0)} &= \frac{1}{y} \quad , \quad A_t = \sqrt{2} q y \quad . \end{aligned} \quad (14)$$

The asymptotic flat region is now at $y = 0$, the curvature singularity is at $y = \infty$, while the two horizons are at $y_{1,2} = q^{-2}(m \pm \sqrt{m^2 - q^2})$. In the following, we will consider the extremal case, which corresponds to $m = q$ and hence to $y_1 = y_2 = q^{-1}$. By setting

$$y = y_1 \frac{a^2}{r + a^2} = \frac{1}{q} \frac{a^2}{r + a^2} \quad (15)$$

the solution (14) can be turned into

$$\begin{aligned} ds_2^2 &= -\frac{dt^2}{(1 + \frac{a^2}{r})^2} + \frac{1}{Q^2} \frac{dr^2}{r^2} \quad , \\ e^{-2(\phi-\phi_0)} &= \frac{q}{a^2} (r + a^2) \quad , \quad A_t = \sqrt{2} \frac{a^2}{r + a^2} \quad . \end{aligned} \quad (16)$$

The two-dimensional (t, r) part of (12) thus agrees with the charged black-hole solution (16) provided that the parameters a, Q and q appearing in (16) are related to the charges

Q_1, Q_5 and Q_{KK} as follows:

$$a^2 = Q_1 \quad , \quad Q^2 = \frac{1}{Q_5 Q_{KK}} \quad , \quad qe^{-2\phi_0} = \frac{Q_1}{\sqrt{Q_5 Q_{KK}}} \quad . \quad (17)$$

The field strength of the gauge field in (16) agrees with $H_{rty} = \partial_r B_{ty}$ given in (3), up to a constant.

The entropy of the charged two-dimensional black-hole (14) was computed in [3, 10] using thermodynamic methods. For an extremal black-hole, it is given by

$$S = \frac{1}{4G_2} e^{-2\phi_0} m \quad . \quad (18)$$

Here, G_2 denotes the two-dimensional Newton's constant, which was set to $G_2 = 1/(16\pi)$ in [3, 10]. Inspection of (12) shows that G_2 is related to Newton's constant G_4 in four dimensions by

$$\frac{1}{G_2} = \frac{V}{G_4} \quad , \quad V = 4\pi Q_5 Q_{KK} \quad , \quad (19)$$

where V denotes the area of the two-sphere with radius $\sqrt{Q_5 Q_{KK}}$. Inserting (19) as well as (17) into the entropy formula (18) yields

$$S = \frac{1}{4G_2} e^{-2\phi_0} m = \frac{\pi}{G_4} Q_1 \sqrt{Q_5 Q_{KK}} \quad , \quad (20)$$

in agreement with the macroscopic formula (5) of the four-dimensional black-hole.

We note here that we expect each of the decoupled two-dimensional geometries of (12) to be described by an exact CFT. We refer to [11, 12, 13, 14] for a discussion of the CFT description of the two-dimensional charged black-hole (13).

We would now like to argue that there is a sequence of S and T - S - T duality transformations which can be used to shift the constant part of the harmonic functions H_5 and H_{KK} . Consider dimensionally reducing the ten-dimensional line element (2) in the t, y, x_6, x_7, x_8, x_9 directions, that is consider compactifying it on a six-torus with Lorentzian signature. The resulting four-dimensional metric in the Einstein frame and the four dimensional dilaton read

$$ds_4^2 = H_{KK} dx_i^2 + \frac{1}{H_{KK}} (dz + A_i dx^i)^2 \quad , \quad e^{-2\phi} = \frac{1}{H_5} \quad . \quad (21)$$

The antisymmetric tensor field H_{zij} given in (3) can be dualised to an axion field b , which can then be combined with $e^{-2\phi}$ into $S_{\pm} = b \pm e^{-2\phi}$ [15]. $S_{\pm}$ can undergo $SL(2, R)$ transformations. The form of the line element (21) makes it clear that, in the case that $H_1 = H_{KK} = 1$, these $SL(2, R)$ transformations can be used to shift away the constant part of H_5 . Similarly, in the case that $H_1 = H_5 = 1$, there exists [15] a sequence of T - S - T

duality transformations which can be used to remove the constant term in H_{KK} . In view of that, we expect that also in the case where H_1, H_5 and H_{KK} are all non-constant, there is a certain sequence of S and T - S - T duality transformations which is capable of removing the constant terms in H_5 and H_{KK} . If so, the solutions (2) with and without constant terms in H_5 and H_{KK} would be dual to one another.

Acknowledgements

This work is supported by the European Commission TMR programme ERBFMRX-CT96-0045.

References

- [1] S. Hyun, *U-duality between Three and Higher Dimensional Black-Holes*, hep-th/9704005.
- [2] E. Teo, *Statistical Entropy of Charged Two-dimensional Black-Holes*, hep-th/9803064.
- [3] M. D. McGuigan, C. R. Nappi and S. A. Yost, *Nucl. Phys.* **B375** (1992) 421, hep-th/9111038.
- [4] G. W. Gibbons and M. J. Perry, *Int. J. Mod. Phys.* **D1** (1992) 335, hep-th/9204090.
- [5] M. Cvetič and A. Tseytlin, *Phys.. Lett.* **B366** (1996) 95, hep-th/9510097.
- [6] M. Cvetič and D. Youm, *Phys. Rev.* **D53** (1996) 584, hep-th/9507090.
- [7] K. Sfetsos and K. Skenderis, *Microscopic Derivation of the Bekenstein-Hawking Entropy Formula for Non-Extremal Black Holes*, hep-th/9711138.
- [8] V. Balasubramanian and F. Larsen, *Near Horizon Geometry and Black-Holes in Four Dimensions*, hep-th/9802198.
- [9] M. Bañados, C. Teitelboim and J. Zanelli, *Phys. Rev. Lett.* **69** (1992) 1849, hep-th/9204099.
- [10] C. R. Nappi and A. Pasquinucci, *Mod. Phys. Lett.* **A7** (1992) 3337, gr-qc/9208002.
- [11] N. Ishibashi, M. Li and A. R. Steif, *Phys. Rev. Lett.* **67** (1991) 3336.
- [12] J. H. Horne and G. T. Horowitz, *Nucl. Phys.* **B368** (1992) 444, hep-th/9108001.
- [13] C. V. Johnson, *Phys. Rev.* **D50** (1994) 4032, hep-th/9403192.
- [14] K. Sfetsos and A. A. Tseytlin, *Nucl. Phys.* **B427** (1994) 245, hep-th/9404063.
- [15] I. Bakas, *Phys. Lett.* **B343** (1995) 103, hep-th/9410104.